2018 November 8th

Dr. Elizabeth Soto Cabrera – Hospital Angeles Tijuana, Mexico

■ Criminal Fraud →

Dr. Elizabeth Soto Cabrera:

SEP (Somatosensory Evoked Potential) – Fraudulated Medical Test Dr. Elizabeth Cabrera

This is a clear example of a doctor falsifying medical reports around data to try and hide the disease pathology around MS (specifically to try and hide the neurodegeneration in the spinal column). In this case the doctor does a Somatosensory Evoked Potential. Its checks the nerve Jan conduction rate over the spinal column (it checks for lesions or neurodegeneration in the spinal column).



The spinal column when this test was done looks like this:

Showing damage to the upper spinal column that would effect movement of limbs (hands and feet).

Narendra

Jana

In the medical results the doctor simply erases all the values that show the reduction in mobility caused by the disease, namely the values for amplitudes.

Id de archivo: NOV0718 Visita: 2018-11-08 08 Nov 18 12:21 Narendra NEURODIAGNOSTICO SCA Nicolet VikingSelect Jana **VEP Plus** Jana, Narendra #1 LED Goggles Registrar 12:21:29 p.m. Det. Vel.: 4 Cambiar: 1.9 Hz · Lado: Izqui. Dur: 5ms Estimulador 100 N Media: Des. Montaje Amplificador Marcadores Estimulador Medida General Notas Ojo izquierdo Texto Lat Lat Lat Amp PP N75 P100 N145 75-100
 ms
 ms
 ms
 uV

 MO
 84.0
 134
 200
 15.8
Ojo derecho
 Texto
 Lat
 Lat
 Lat
 Amp PP

 N75
 P100
 N145
 75-100

 ms
 ms
 uV
 200
MO 83.8 138 213 7.19 Ambos ojos
 Texto
 Lat
 Lat
 Lat
 Amp PP

 N75
 P100
 N145
 75-100

 ms
 ms
 ms
 uV
MO 90.0 128 196 17.3 Medidas interoculares MOEzM Texto Dif. lat Rel amp. PP LT-RT LT-RT ms 44 MO 4.50 45.5 Σ QIL MOFzMF 25ms 20uV 1 20.1 T Edición: Des.

But the graphs show the clear effects that would cause a difficulty in movement due to neurodegeneration, inflammation, or lesions in the spinal column:

The first graph checks for visual evoked potentials (VEP), it does acknowledge optic neuropathy and isn't falsified.



In the above graph the difference in amplitude (the difference in difficulty of movement) is readily apparent, the sharp peaks in the lower 4 graphs are in contrast to the clear diminished amplitudes in the upper graph. I have better movement in the right limbs than the left. The results from this graph are falsified. The graphs are also better formed in the lower graph then the upper, indicating neurodegeneration effecting the left hemisphere.



The values missing the last graph indicate that the graphs was so poor that the machine couldn't measure it, the LP-P37 results for the left limb and the N41P37 values for the left limb. The SUBCO value is also deleted or couldn't be measure due to clinical effect. The graphs for the right hemisphere are also better formed then the left indication neurodegeneration that effects the left hemisphere. They are clear abnormalities in a patient with MS. The doctor then falsifies her report based on falsified results:

Narendra DRA. ELIZABETH SOTO CABRERA NeuroDiagnóstico SCA Jana NEURÓLOGA Y NEUROFISIOLOGÍA CLÍNICA Visual Evoked Potentials, Upper and Lower Limbs Cédula Profesional Somatonsensory Evoked Potentials Patient Patient Narendra Jana Date 11/08/2018 Age 34 yo Sex Male Handedness Right Diagnosis Dr. Gutiérrez Sent Perform Dra. Elizabeth Certificada por el study Soto Cabrera Technique FINDINGS: Visual Evoked Potentials (VEP) Monocular stimulation by stroboscopic light with goggles and placen record electrodes in Oz. We obtained integration of component P100 on the left side with latency of 134 ms and on the right side, latency of 138 ms, with bilateral stimulation we obtained a latency of 128 ms (normal in under 132ms with goggle stimulation). Interocular latency difference is 4ms (normal VN <10 ms). Amplitude in right side is less tan 50% comparing to left side (7.19 uV vs 15.8 uV) wich is significative. Lower Limbs Somatosensory Evoked Potentials (SEP) We stimulated Posterior Tibial Nerve bilaterally. Record electrodes were placed in Cz, C5, L1 and popliteal fossa bilaterally. We applied 250 stimulus with 3.1 Hz, with respective replication in each side. We obtained integration of component in Popliteal fossa (HP), Lumbar and Cortical. We found normal latencies bilaterally in every component. There is no side to side significative difference. Upper Limbs Somatosensory Evoked Potentials (SEP) facebook.com/neurologaelizabethsot 🔯 elizabethsotoneurologa@gmail.com We stimulated Median Nerve bilaterally. Record electrodes were placed in C3, C4, CV and Erb bilaterally. We applied 250 stimulus with 3.1 Hz, with respective replication in each side. We obtained integration of component Erb, Twitter @NeurologaESoto CV, SC and cortical. We found normal latencies bilaterally in every component. There is no side to side significative difference. Paseo de los Héroes No. 10999 7mo. Piso, Consultorio 703 Zona Río Tijuana, C.P. 22010, Tijuana, B.C. Teléfonos: (664) 635.1871 y (664) 635.1800 ext 6703 0



Jana

The SEP of the upper and lower limbs have clear differences in amplitude and possibly latency (difficult to determine due to malformed graphs) effecting movement and determined by both the MRI images and the graphs in the tests.

This medical diagnostic and report stands out as one of the easiest examples of demonstrating an attempt at falsifying a diagnostic and shows a clear case of fraudulence with an attempt to hide a condition to further neurodegeneration or to hide harm from past negligence.

Narendra

Jana